

LETTERS

HEALTH COMMUNICATION, INTELLIGENCE, AND HEALTH DIFFERENTIALS

In a recent editorial, Freimuth and Quinn¹ discussed the contribution that health communication might make to eliminating health disparities. The mechanisms underlying the recently suggested role of intelligence (denoted here by IQ) in the etiology of several important health outcomes are pertinent to this debate.

In long-term follow-up studies, persons with low scores on written IQ tests administered in childhood and early adulthood have been found to have elevated rates of all-cause mortality,² coronary heart disease,³ some cancers,⁴ and psychiatric disorders⁵ than their higher-scoring counterparts in later life. For the most examined health outcome—all-cause mortality—these effects are strong (exceeding those of other psychological characteristics, such as personality disposition, and on a par with the effects of established risk factors such as raised blood pressure, social disadvantage, and obesity), incremental (apparent across the full range of IQ scores, not just confined to below-average scores), consistent across a number of study populations and research groups, and seemingly independent of childhood social circumstances.⁶

IQ reflects an individual's ability to learn, reason, and solve problems.⁷ Therefore, plausible mechanisms underlying these observations that have particular relevance to health communications—defined by Freimuth and Quinn¹ as the study of methods that influence decisions that enhance health—concern the conceptualization of self-health care as a set of cognitive tasks. These tasks include the optimal interpretation of health prevention messages and, among persons with existing chronic illness, improved disease management.⁶

In terms of the former, Scottish children who scored high on tests of intelligence in 1932 were more likely than others to stop smoking as adults in the 1960s and 1970s.⁸ This observation may be attributable to the differential interpretation of antismoking advice that appeared for the first time during this period. With regard to disease management, the daily management of one's illness is cognitively demanding; it requires ongoing learning and independent decisionmaking. Persons with low levels of education or health literacy, both of which are related to IQ,⁷ are less likely to appreciate when their conditions require medical attention, to be aware of the appropriate actions to take when they do realize they need attention, and to correctly comprehend instructions for self-medication.⁹ They may also fail to seek medical advice promptly after a significant illness episode (e.g., myocardial infarction) and receive treatment at a facility most appropriate to their clinical requirements.⁹

Health care advice, treatment protocols, and disease prevention information may exceed the intellectual capacity of some people.^{9,10} If this is the case, and since highly simplified advice might substantially reduce the intended impact of the advice, proactive involvement of health care providers in the provider–client interaction, as advocated by Freimuth and Quinn,¹ might reduce health differentials. ■

G. David Batty, PhD, MSc
Ian J. Deary, PhD, FRCPE

About the Authors

The authors are with the Department of Psychology, University of Edinburgh, Edinburgh, Scotland. G. David Batty is also with the Medical Research Council Social and Public Health Sciences Unit, University of Glasgow, Glasgow, Scotland.

Requests for reprints should be sent to G. David Batty, MRC Social and Public Health Sciences Unit, University of Glasgow, 4 Lilybank Gardens, Glasgow, UK, G12 8RZ (e-mail: david-b@msoc.mrc.gla.ac.uk).

doi:10.2105/AJPH.2005.063370

Acknowledgments

G. David Batty is the recipient of a Wellcome Advanced Training Fellowship (071954/Z/03/Z); Ian Deary holds a Royal Society-Wolfson Research Merit Award.

References

1. Freimuth VS, Quinn SC. The contributions of health communication to eliminating health disparities. *Am J Public Health*. 2004;94:2053–2055.
2. Whalley LJ, Deary IJ. Longitudinal cohort study of childhood IQ and survival up to age 76. *BMJ*. 2001;322:819.
3. Batty GD, Mortensen E, Nybo Andersen A-M, Osler M. Childhood intelligence in relation to adult coronary heart disease and stroke risk: evidence from a Danish birth cohort study. *Paediatr Perinat Epidemiol*. In press.
4. Hart CL, Taylor MD, Davey Smith G, et al. Childhood IQ, social class, deprivation, and their relationships with mortality and morbidity risk in later life: prospective observational study linking the Scottish Mental Survey 1932 and the Midspan studies. *Psychosom Med*. 2003;65:877–883.
5. Batty GD, Mortensen EL, Osler M. Childhood intelligence in relation to later psychiatric disorder: evidence from a Danish birth cohort study. *Br J Psychiatry*. In press.
6. Batty GD, Deary IJ. Early life intelligence and adult health. *BMJ*. 2004;329:585–586.
7. Neisser U, Boodoo G, Bouchard TJ Jr, et al. Intelligence: knowns and unknowns. *Am Psychol*. 1996;51:77–101.
8. Taylor MD, Hart CL, Davey Smith G, et al. Childhood mental ability and smoking cessation in adulthood: prospective observational study linking the Scottish Mental Survey 1932 and the Midspan studies. *J Epidemiol Community Health*. 2003;57:464–465.
9. Gottfredson LS, Deary IJ. Intelligence predicts health and longevity, but why? *Curr Dir Psychol Sci*. 2004;13:1–5.
10. Gottfredson LS. Intelligence: is it the epidemiologists' elusive "fundamental cause" of social class inequalities in health? *J Pers Soc Psychol*. 2004;86:174–199.

Letters to the editor referring to a recent Journal article are encouraged up to 3 months after the article's appearance. By submitting a letter to the editor, the author gives permission for its publication in the Journal. Letters should not duplicate material being published or submitted elsewhere. The editors reserve the right to edit and abridge letters and to publish responses.

Text is limited to 400 words and 10 references. Submit online at www.ajph.org for immediate Web posting, or at submit.ajph.org for later print publication. Online responses are automatically considered for print publication. Queries should be addressed to the department editor, Jennifer A. Ellis, PhD, at jae33@columbia.edu.